



FSN-EN 0003

TOXICITY OF DRILLING FLUID XP-07 ON MICROBIAL LOAD AND SURVIVAL OF *OREOCHROMIS NILOTICUS* FINGERLINGS

*VINCENT-AKPU, I. F. AND SIKOKI, F. D.

Department of Animal and Environmental Biology,
Faculty of Science, University of Port Harcourt, Rivers State, Nigeria

Copyright 2010, Fisheries Society of Nigeria.

This paper was prepared for presentation at the 25th Annual International Conference and Exhibition in Administrative Staff College of Nigeria (ASCON), Topo-Badagry, Lagos, Nigeria, 25th – 29th October, 2010.

This paper was selected for presentation by an FISON Program Committee following review of information contained in an abstract submitted by the author(s). Contents of the paper, as presented, have not been reviewed by the Fisheries Society of Nigeria and are subject to correction by the author(s). The material, as presented, does not necessarily reflect any position of the Fisheries Society of Nigeria, its officers, or members. Papers presented at FISON meetings are subject to publication review by Editorial Committees of the Fisheries Society of Nigeria. Electronic reproduction, distribution, or storage of any part of this paper for commercial purposes without the written consent of the Fisheries Society of Nigeria is prohibited. Permission to reproduce in print is restricted to an abstract of not more than 300 words; illustrations may not be copied. The abstract must contain conspicuous acknowledgement of where and by whom the paper was presented. Write Librarian, Fisheries Society of Nigeria (FISON), P. O. Box 2607 Apapa, Lagos.

ABSTRACT

Static bioassays were carried out in the laboratory for 96h to investigate the effect of Parateq a synthetic base mud on the fingerling of *Oreochromis niloticus* and their microbial population. The fingerlings were exposed to different concentrations (0, 1000, 3000, 5000 and 9000Mg/l) of drilling fluid. Fish mortality data were recorded at 24, 48, 72 and 96h and analysed using Finney's Probit Analysis method to calculate LC_{50} values (concentration of drilling fluid in water that will kill 50% of the fish population in 96 hours). The 96h LC_{50} was 2210Mg/l. The mortality increased with increase in concentration, while, the lethal times decreased as concentration increased. The bacterial biomass ranged from 1.59×10^8 to 6.0×10^6 CFU/ml while fungal biomass ranged from 1.0×10^3 to 2.1×10^4 CFU/ml. Results revealed that there was an increase in fungal biomass with increase in mud concentration and a decrease in bacterial biomass with increase in mud concentration.

INTRODUCTION

In West Africa, Nigeria (Niger Delta) a lot of oil and gas exploration and development activities are carried out

using drilling fluids. Drilling fluids are suspension of solid in liquid emulsion and/or dissolved materials with chemical additives which are employed during exploration to remove cuttings. Two types of fluids normally use in drilling operations are water based fluids and non-aqueous drilling fluids (OGP, 2003). Drilling fluid when discharged untreated into the environment are capable of interfering with the normal functions of the organisms (Saasen *et al.*, 2001; Wills, 2000). Drilling waste may kill marine life, smother or suffocate it with plume of suspended particles. This will result in potential damage to fish and invertebrate population and alteration in feeding and spawning grounds (Neff *et al.*, 2000; Soegianto *et al.*, 2008). Rhodes and Hendricks (1990) discovered that the components of drilling fluids inhibit the growth of microbial communities that are important in some biological cycles present in the ecosystem which may affect the productivity of such system.

In aquatic ecosystems, Fish are located at the top of the food chain and are highly visible resources. They are known to be affected by pollutants due to their being in direct contact with water via their gills and body surface. Fish and fishery products are generally regarded as high risk commodity in respect of pathogen contents, natural toxins and possible contaminants and alterations (Yousuf *et al.*, 2008). The estimation of bacterial numbers in fish is frequently used to retrospectively assess microbial quality or to assess the presumptive safety of the product (Okonko *et al.*, 2008). When microbial growth and actions are affected by the presence of contaminants in the aquatic system, the fishes are exposed to

harmful organisms which are subsequently passed on to man.

Ekpo and Uzegbu, (2004) studied the effect of drilling additives (chromelignsulphonate) on *Pachymelania fusca* and *Tympanotonus fuscatus* and observed an increase in the microbial biomass with increase in concentration of drilling additives. However, little information exists on effect of drilling fluid and its additives on microbial load of fish. This study aims to determine any significant changes in the bacterial and fungal biomass in *Oreochromis niloticus* when exposed to Parateq a commonly used drilling fluid in the Niger Delta wetlands of Nigeria.

MATERIALS AND METHODS

Fingerlings (8.45cm) of *Oreochromis niloticus* were obtained from the African Regional Aquaculture Centre (ARAC), Aluu, Rivers State - Nigeria.

In the laboratory, the fish were transferred immediately to the holding tanks [120cm x 120cm x 120cm]. The holding tanks were aerated, cleaned and the water renewed daily (Reish and Oshida, 1986). Fish were fed twice daily with Nigeria Institution of Oceanography and Marine Research feed (NIOMR) (35% protein).

Drilling fluid was obtained from Bariods Nigeria Ltd and prepared in advance. The fluid was stored in a 5 litre bucket at approximately 5°C. Immediately before the fluid was used, it was mixed in a storage bucket with a mixer at 3450 rpm for ½ hour to ensure sample homogeneity.

A complete randomised designed was used and the tests were conducted in the laboratory under room temperature using static non-renewal bioassay (Ca1/EPA, 2004). The solutions (0000Mg/l, 1000Mg/l, 3000Mg/l, 5000Mg/l and 9000Mg/l) were thoroughly mixed for 20 minutes and allowed to settle for 1 hour before the introduction of fish (Reish and

Oshida, 1986, FAO, 1987).). The drilling fluid settled into a thin layer covering the bottom of the test chambers.

One hour after the preparation of the test solutions, ten healthy organisms were randomly selected and placed in the tanks in 3 replicates and covered with a net that was fastened with a rubber band to prevent fish escape. The fish were not fed during the 96hrs bioassay. Water quality parameters (Temperature, Alkalinity, pH and Dissolved Oxygen) of the test tanks were monitored daily for the replicates using standard methods (APHA, 1998).

Mortality was used as a measure of toxicity in terms of 96 hrs LC₅₀. The numbers of dead and active fishes were recorded. A fish is presumed dead and removed immediately when there are no opercula or body movement after gentle prodding.

At the end of the 96 hours, the total viable count (TVC) of heterotrophic bacterial and fungal population on the fish body was determined using nutrient agar (APHA, 1998). The average counts of colony forming unit (CFU /ml) were calculated and recorded.


RESULTS AND DISCUSSION

The various physicochemical parameters (Temperature - 28.15 ± 0.65 ; pH - 6.61 ± 0.31 ; Alkalinity - 23.38 ± 0.019 ; Dissolved Oxygen - 3.83 ± 0.28) did not vary significantly ($p > 0.05$) from those of the control except a slight decrease in the Dissolved oxygen (DO) which Ifeadi *et al* (1985) attributed to the effect of oil based fluid.

No mortality was observed in the control, however, in the other treatments, mortality increased with increase in drilling fluid concentration over time. At 24h, no death was recorded at the lowest concentration of Parateq. However, as the time of exposure increased, more deaths were recorded. At the end of 96h, the result showed 26, 33, 73 and 80% mortality recorded in 1000, 3000, 5000

and 9000 mg/l respectively (Fig. 1). The LT_{50} also decrease with increase in

concentration.



The results indicated that as the concentration increased, more deaths were recorded and as the time increased the LC_{50} reduced (Table 1). The toxicity of Drilling fluids has been attributed to their hydrocarbon content (Neff *et al.*, 2000; Tagatz *et al.*, 1985). Low Hydrocarbon content in the drilling fluid can be responsible for the low toxicity and also Synthetic based fluid does not readily disperse in water (Wills, 2000). Parrish and Ward (1988) recorded LC_{50} range of 0.45-39% using laboratory prepared drilling fluid on mysid and posited that the increased toxicity was due to the presences of hydrocarbon. The 96hrs LC_{50} value of the static bioassay with Parateq was 2210 mg/l (1% = 1000 mg/l), this LC_{50} was relatively low. This is probably because the test was not renewed daily during the 96h bioassay period (Neff *et al.*, 1981).

Table 2 shows the heterotrophic fungal and bacterial counts of *O. niloticus* exposed to Parateq. The total fungal counts ranged from 1.0×10^3 to 2.1×10^4 CFU/ml where the highest counts were

found in fish exposed to 9000Mg/L. The fish in control tanks and in concentration 1000Mg/L had total viable count of 1.0×10^3 . While in the other concentrations, the cells increased from the lowest to the highest concentration. The bacterial count decreased with increase in Parateq concentration. The fish in control had bacterial counts of 1.59×10^8 CFU/ml while those in highest concentration (9000Mg/L) had a bacterial count of 6.0×10^6 CFU/ml. The total counts for all the samples examined in this study were generally high exceeding the FAO/WHO Standard limits of 1.0×10^2 CFU/ml that can be present in fish (ICMSF, 1986). This indicated that there is presence of harmful microorganisms on the fish body. Similar trend of bacterial growth was reported by Ekpo and Uzegbu, (2004) where the coliform counts ranged between 5.4×10^2 and 8.5×10^5 for shrimp and 5×10^2 to 4.4×10^4 cells for prawn when exposed to drilling mud additives. They further stated that when the total count reaches 10^6 CFU per gram or millilitre, the product is assumed to be, or nearing, spoilage. As noted by earlier workers

(Smith *et al.*, 1992), micro organisms are usually the normal flora on the body of the fish. Their growth and multiplication are either encouraged or inhibited by toxicants. In this study, the growth of the bacterial population was inhibited while the growth of the fungal population was encouraged. Nweke and Okpokwasili (2003) observed that 1% drilling fluid waste elicited differential toxic effects on four bacterial isolates from the mangrove swamps in the Niger Delta. This they attributed to the natural ability of some bacteria to biodegrade the base oil. When bacterial growth and action are hindered by the presence of drilling fluid, fishes are exposed to harmful materials.

CONCLUSION

Despite the importance and sensitive nature of the Niger Delta, the area is replete with oil exploration and exploitation activities, which involve the use of drilling fluids. Most of the cuttings produced contain considerable proportion of drilling fluids which are dumped at the well site or deposited into the water. Drilling fluids elicited differential effects on the microbial population on the fish body. This has been attributed to the natural ability of some bacteria to biodegrade the base fluid. Thus micro organisms can increase or decrease in number depending on the effect of base fluid on the species.

REFERENCES

- APHA, 1998. Standard methods for the examination of water and wastewater, 20th ed. APHA-AWWA-WPCF. Washington D.C. 1220.
- Cal/EPA, 2004. Overview of fresh water and marine Toxicity Tests: A Technical Tool for Ecological Risk Assessment. California. Environmental Health Hazard Assessment and Dept. of Environmental Toxicology, University of California. Davis. 147.
- Ekpo A. E. and Uzegbu M. L., 2004. The effect of petroleum drilling mud additives (Chromelignsulphonate) on the microbiology and survival periwinkle (*Pachymelania spp.* and *Tympanotonus sp.*) *World Journal of Microbiology and Biotechnology*. 20: (6) 559-562.
- FAO, 1987. Bases for selecting biological tests to evaluate marine environment. Manual of methods in aquatic research. 4. 4
- ICMSF (International Commission on microbiological Specifications for Foods). 1986. Microorganisms in foods 2-Sampling for microbiological analysis: principles and specific applications. Ontario, Canada, University of Toronto Press (2nd edition)
- Ifeadi, C. N., Nwankwo, J. N., Ekaluo, A. B. and Orubima, I. I. 1985. Treatment and disposal of drilling muds and cuttings in the Nigeria Petroleum Industry. The Petroleum Industry and the Nigeria Environment. Proceeding of 1985 seminar Kaduna Nigeria. 55-80.
- OGP (Oil and Gas Producer), 2003. Environmental Aspects of the Use and Disposal Of Non-Aqueous Drilling Fluids Associated with Offshore Oil and Gas Operations. Report No. 342.
- Okonko, I. O., Ogunnusi, T. A., Ogunjobi, A. A., Adedeji, A. O., Adejoye, O. D., Babalola, E. T. and Ogun A. A. 2008. Microbial studies on frozen shrimps processed in Ibadan and Lagos, Nigeria. *Scientific Research and Essay*. 3 (11), 537-546.
- Neff, J. M., McKelvie, S. and Ayers, R. C. Jr. 2000. A literature review of environmental impacts of synthetic based drilling. Report to U.S. Dept of the interior minerals

- management service. Gulf of Mexico OC office New Orleans, LA. 154.
- Neff, J. M., Carr, R. S. and McCulloch, W. L. 1981. Acute toxicity of a used chrome lignosulphonate drilling mud to several species of marine invertebrate. *Marine Environ Res.* 4. 251 – 266.
- Nweke, G. C. and Okpokwasili, I. C. 2003. Drilling fluid base oil biodegradation potential of a soil. *Staphylococcus species. Africa J. Biotech.* 2(9) 293 – 295.
- Parrish, P. R. and Ward, T. W. 1988. Variability of the toxicity of drilling fluids to mysids (*Mysidopsis bahia*). In: chemical and biological characterization of sludges, sediments, dredge spoils and drilling muds. ASTM STP 976. Philadelphia. 326 – 333.
- Reish, D. J. and Oshida, P. S. 1986. Manual of Methods in Aquatic Environmental Research. Part 10. Short-term static bioassays. FAO fish. Tech Pap (247) 62.
- Rhodes, A. N., and Hendricks, C. W. 1990. A continuous flow method for measuring effect of chemicals on soil nitrification. Toxicity Assessment. 5. 77- 89.
- Saasen, A., Berntsen, A., Loklingholm, M. G., Igeltjom, G. and Asnes, K. 2001. The effect of drilling fluid base-oil properties on occupational Hygiene and the Marine Environment. SPE drilling and completion 16 (3) 150 – 153.
- Smith, G. A., Nickels, J. S. Bobbie, R. J. Rich, N. L. E and White, D. C. 1992. Effects of oil and gas well drilling fluids on the biomass and community structure of micro biota that colonize sand and running waters. *Altantica.* 5(2). 39.
- Soegianto, A., Bambang, I., Mochammad, A. 2008. Toxicity of drilling waste and its impact on gill structure of post larvae of tiger prawn (*Penaeus monodon*) *Global Journal of Environmental Resource* 2(1): 36 - 41.
- Tagatz, M. E., Plaia, G. R. and Deans, C. H. 1985. Response of macrobenthos colonizing estuarine sediments contaminated with drilling mud containing diesel oil. *Bull. Environ. Contam. Toxicol.* 35: 112-120.
- Wills, J. 2000. Muddied waters. A survey of oil field drilling wastes and disposal techniques to reduce the ecological impact of sea dumping. Sakhalin Environmental Watch. 139.
- Yousuf A. H. M., Ahmed, K. Yeasmin S., Ahsan N., Rahman M. and Islam M. 2008. Prevalence of microbial load in shrimp *Penaeus monodon* and prawn, *Macrobrachium rosebergii* from Bangladesh. *World Journal of Agriculture Science* 4 (S): 852 – 855.

Table 1: Median Lethal concentrations (LC₅₀) of Parateq on *O. niloticus* in 96 hours

Time (hours)	LC ₅₀ (Mg/L)
48	3310
72	2710
96	2210

Table 2: Viable count of colonies from the fungal and bacterial isolates

Concentrations (Mg/L)	Fungal Counts(CFU /ml)	Bacterial Counts(CFU /ml)
0 000	1.00 x 10 ³ ± 141.4	1.59 x 10 ⁸ ± 79.2
1000	1.50 x 10 ³ ± 282.8	1.55 x 10 ⁸ ± 8.50
3000	1.00 x 10 ³ ± 282.8	1.12 x 10 ⁸ ± 15.6
5 000	1.70 x 10 ³ ± 282.8	2.70 x 10 ⁷ ± 8.5
9000	2.10 x 10 ³ ± 424.3	6.00 x 10 ⁶ ± 1.4